


# Coronal signatures of flares and CMEs

Dr. Karin Muglach  
NASA/GSFC and CUA

SW-REDI 2016






# Large Scale Structures Near the Solar Surface

two kinds of measurement to collect information about the Sun:

**Remote Sensing** and **In-situ Measurement**

# Key for remote sensing of the sun (and stars): Solar Spectrum


Solar Spectrum





complete solar spectrum  
and  
EUV part of solar spectrum

# Key for remote sensing of the sun (and stars): Solar Spectrum

*True-Color Irradiance Spectrum 392 to 692 nm from Kitt Peak Residual Irradiance Atlas (Kurucz 2005)*





# Key for understanding solar activity: the solar magnetic field



Global magnetic field (extrapolation): 3d structure

Line-of-sight full disk magnetogram: 2d cut at photosphere

# Key for understanding solar activity: the solar magnetic field



Full disk white light image (SDO), full disk line-of-sight magnetogram (SDO)

# Key for understanding solar activity: the solar magnetic field

Active Region evolution in white light and magnetogram (SDO).

# Key for understanding solar activity: the solar magnetic field

If we just have white light images and magnetograms:

Q: How are the polarities connected?

# Key for understanding solar activity: the solar magnetic field



If we just have white light images and magnetograms:

Q: How are the polarities connected?

A1: extrapolation


A2: corona images: outline (some) of the magnetic field  
connectivity!

# Key for understanding solar activity: the solar magnetic field




Full disk magnetogram and 171 image (SDO)

# Key for understanding solar activity: the solar magnetic field



Full disk image  
in H alpha  
(BBSO):  
filaments seen  
as dark  
absorption  
structures

# Key for understanding solar activity: the solar magnetic field




High resolution  
image in H  
alpha (Dutch  
Open  
Telescope)  
filaments seen  
as dark  
absorption  
structures

# Key for understanding solar activity: the solar magnetic field


Example of filaments:

- Quiescent filament in high spatial resolution (Hinode SOT)
- Filament eruption (SDO, composite)

# SIMPLE (!! ) cartoon of active region magnetic field



# SIMPLE (!! ) cartoon of filament magnetic field



magnetic field lines almost  
parallel to PIL

# Key for understanding solar activity: the solar magnetic field

Notes on filaments:

- Filament: on-disk structure (seen in absorption)  
Prominence: same structure off limb (seen in emission)
- Best wavelengths: H alpha, He II 304, Fe XII 195 A (AIA, STEREO)
- All filaments have a PIL
- But not all PILs are filaments!
- Caution: full disk magnetograms give only the line-of-sight magnetic field – projection effects near the solar limb!

# Solar Eruptions: Flares and CMEs

- Energy is stored in the solar magnetic field (active regions and filaments): accumulated over a long period of time – days, weeks, months
- Energy is released in eruptions (flares, CMEs): in a short time scale (minutes, hours)


Magnetic energy is converted to thermal energy (and radiative energy) and kinetic energy (e.g. mass motion in CMEs and SEPs)

# Solar Eruptions: Flares and CMEs


Solar Flares: Event that releases X-rays

X-ray monitor on-board GOES spacecraft (in Earth orbit), full disk monitor (no spatial information of location of flare on the sun)

larger events radiate also in other wavelengths especially in UV, EUV (and radio) → use SDO/AIA images to determine location!



# Solar Eruptions: Flares and CMEs



one possible  
scenario for an  
eruption:

- reconnection at the x-point (energy release)
- CME escapes upward, field-lines open up
- Post-eruptive loops appear below x-point (additional heating)

# Solar Eruptions: Flares and CMEs

**Caution:** the real sun is more complicated compared to the cartoon – e.g. magnetic field is a

## 3d structure

- some eruptions show no/very little X-ray signature (particularly filament eruptions)
- some flares have no CMEs

# Large scale structures in the corona

- Images: SDO AIA 193 Å, STEREO EUVI 195 Å  
(filter contains Fe XII 195 Å line,  $T \sim 1.5$  MK)
- Line-of-sight magnetograms: polarity inversion line (PIL)
- **Active Regions:** bi-polar, bright (emission), closed magnetic field (field lines perpendicular to PIL)
- **Filaments:** bi-polar, dark (absorption), closed magnetic field (field lines parallel to PIL)
- **Coronal hole:** uni-polar, dark (less dense), open magnetic field

# Coronal signatures of CMEs

- Data to use: SDO AIA, STEREO EUVI (A & B)
- **Brightenings**: flares, post-eruptive arcade (193), arcade footpoints (304, 193)
- **Darkenings**: dimmings (transient coronal holes), dark/absorbing/cool material rising (filament eruption)
- **Off-limb**: opening of closed coronal field lines, AIA 304 emission structure
- Not a signature of eruption: active region loop brightenings, (small) flares


# Coronal signatures of CMEs

Good period to study: SDO 2014-02-18 - 21  
(use AIA 211, 193, 304)

# Coronal signatures of flares and CMEs

Dr. Karin Muglach  
NASA/GSFC and CUA

SW-REDI 2016





# Large Scale Structures Near the Solar Surface

two kinds of measurement to collect information about the Sun:

**Remote Sensing** and **In-situ Measurement**


# Key for remote sensing of the sun (and stars): Solar Spectrum

Solar Spectrum




ar spec

solar sp





# Key for remote sensing of the sun (and stars): Solar Spectrum



*True-Color Irradiance Spectrum 392 to 692 nm from Kitt Peak Residual Irradiance Atlas (Kurucz 2005)*



# Key for understanding solar activity: the solar magnetic field



# Key for understanding solar activity: the solar magnetic field



k lin

# Key for understanding solar activity: the solar magnetic field

Active Region evolution in white light and magnetogram (SDO).

## Key for understanding solar activity: the solar magnetic field

If we just have white light images and magnetograms:

Q: How are the polarities connected?

## Key for understanding solar activity: the solar magnetic field



If we just have white light images and magnetograms:

Q: How are the polarities connected?

A1: extrapolation


A2: corona images: outline (some) of the magnetic field  
connectivity!

## Key for understanding solar activity: the solar magnetic field




ge

## Key for understanding solar activity: the solar magnetic field



Key for understanding solar activity:  
the solar magnetic field




# Key for understanding solar activity: the solar magnetic field

Example of filaments:


- Quiescent filament in high spatial resolution (Hinode SOT)
- Filament eruption (SDO, composite)

# SIMPLE (!!)

cartoon of active region magnetic field



SIMPLE (!! ) cartoon of filament magnetic field



magnetic field lines almost  
parallel to PIL

# Key for understanding solar activity: the solar magnetic field

Notes on filaments:

- Filament: on-disk structure (seen in absorption)  
Prominence: same structure off limb (seen in emission)
- Best wavelengths: H alpha, He II 304, Fe XII 195 A (AIA, STEREO)
- All filaments have a PIL
- But not all PILs are filaments!
- Caution: full disk magnetograms give only the line-of-sight magnetic field – projection effects near the solar limb!

# Solar Eruptions: Flares and CMEs

- Energy is stored in the solar magnetic field (active regions and filaments): accumulated over a long period of time – days, weeks, months
- Energy is released in eruptions (flares, CMEs): in a short time scale (minutes, hours)


Magnetic energy is converted to thermal energy (and radiative energy) and kinetic energy (e.g. mass motion in CMEs and SEPs)


# Solar Eruptions: Flares and CMEs

Solar Flares: Event that releases X-rays

X-ray monitor on-board GOES spacecraft (in Earth orbit), full disk monitor (no spatial information of location of flare on the sun)

larger events radiate also in other wavelengths especially in UV, EUV (and radio) → use SDO/AIA images to determine location!





# Solar Eruptions: Flares and CMEs

**Caution:** the real sun is more complicated compared to the cartoon – e.g. magnetic field is a

## 3d structure

- some eruptions show no/very little X-ray signature (particularly filament eruptions)
- some flares have no CMEs

# Large scale structures in the corona

- Images: SDO AIA 193 Å, STEREO EUVI 195 Å  
(filter contains Fe XII 195 Å line,  $T \sim 1.5$  MK)
- Line-of-sight magnetograms: polarity inversion line (PIL)
- **Active Regions:** bi-polar, bright (emission), closed magnetic field (field lines perpendicular to PIL)
- **Filaments:** bi-polar, dark (absorption), closed magnetic field (field lines parallel to PIL)
- **Coronal hole:** uni-polar, dark (less dense), open magnetic field

# Coronal signatures of CMEs

- Data to use: SDO AIA, STEREO EUVI (A & B)
- **Brightenings**: flares, post-eruptive arcade (193), arcade footpoints (304, 193)
- **Darkenings**: dimmings (transient coronal holes), dark/absorbing/cool material rising (filament eruption)
- **Off-limb**: opening of closed coronal field lines, AIA 304 emission structure
- Not a signature of eruption: active region loop brightenings, (small) flares

# Coronal signatures of CMEs

Good period to study: SDO 2014-02-18 - 21  
(use AIA 211, 193, 304)